Digitized by the Internet Archive
in 2026

https://archive.org/details/journal-of-chemical-thermodynamics_1992_24_contents

THE JOURNAL OF CHEMICAL THERMODYNAMICS

Volume 24, 1992

EDITORS

M. L. McGlashan
Department of Chemistry
University College London
20 Gordon Street
London WC1H 0AJ
U.K.

P. A. G. O'Hare
National Institute of
Standards and Technology
Building 222, Gaithersburg
MD 20899
U.S.A.

Assistant Editor: **M. B. Ewing** (University College London, U.K.)

ADVISORY BOARD

Chairman: G. Somsen (The Netherlands)

E. H. P. Cordfunke (The Netherlands)
G. Della Gatta (Italy)
J. H. Dymond (U.K.)
J. Fuger (Commission of the European Communities)
L. V. Gurvich (U.S.S.R.)
K. N. Marsh (U.S.A.)
G. M. Olofsson (Sweden)
D. Patterson (Canada)
G. Pilcher (U.K.)
J. A. Rard (U.S.A.)
G. M. Schneider (F.R.G.)
M. Sorai (Japan)
W. V. Steele (U.S.A.)
F. L. Swinton (U.K.)
W. Wagner (F.R.G.)
R. D. Weir (Canada)

ACADEMIC PRESS
(Harcourt Brace Jovanovich, Publishers)
London San Diego New York
Boston Sydney Tokyo Toronto

per.
QD
501
.J683

Copyright © 1992 Academic Press Limited

ALL RIGHTS RESERVED

No part of this volume may be reproduced in any form, by photostat, microfilm, or any other means, without written permission from the publishers.

Printed in Great Britain

CONTENTS OF VOLUME 24

Number 1, *January* 1992

Kabo, G. J.; Kozyro, A. A.; Krouk, V. S.; Sevruk, V. M.; Yursha, I. A.; Simirsky, V. V.; Gogolinsky, V. I. Thermodynamic properties of 6-aminohexanoic lactam (ε-caprolactam) 1

Ortega, J.; Gonzalez, E.; Matos, J. S.; Legido, J. L. Thermodynamic properties of (a methyl ester + an *n*-alkane). I. H_m^E and V_m^E for $\{xCH_3(CH_2)_{u-1}CO_2CH_3$ (u = 1 to 6) + $(1-x)CH_3(CH_2)_6CH_3\}$ 15

García de la Fuente, I.; Rodríguez, J. F.; González, J. A.; Cobos, J. C.; Casanova, C. Excess molar volumes of (*n*-nonan-1-ol + *n*-decane or *n*-tetradecane) at the temperatures 298.15 K, 308.15 K, and 318.15 K 23

Bugajewski, Z.; Bylicki, A. (Solid + liquid) phase equilibria for (*N*-methylpyrrolidin-2-one + 2,6-dimethylphenol) 29

Airoldi, C.; Digiampietri, E. A. Thioamide adducts of zinc-group chlorides—a thermochemical study 33

Róg, G.; Kozłowska-Róg, A.; Zakuła, K. Determination of the standard Gibbs free energies of formation of the sodium aluminates by e.m.f. measurements . . 41

Rard, J. A. Isopiestic determination of the osmotic and activity coefficients of $\{(1-y)H_2SO_4 + yNa_2SO_4\}$(aq) at the temperature 298.15 K. II. Results for y = (0.12471, 0.24962, and 0.37439) 45

Lloyd, M. J.; Wormald, C. J. Excess molar enthalpies of $\{xCH_3)_2CO + (1-x)C_6H_6\}$ at temperatures from 494.5 K to 577.4 K at the pressure 5.68 MPa 67

Ott, J. B.; Sipowska, J. T.; Woolley, A. T.; Izatt, R. M. Excess enthalpies for (propane + butan-1-ol) at the temperatures 298.15 K and 323.15 K and the pressures 5 MPa and 15 MPa, and at 348.15 K and 363.15 K and 5 MPa, 10 MPa, and 15 MPa . 75

Sandhu, J. S.; Singh, A. Excess molar volumes of (acrylonitrile + an *n*-alkan-1-ol} at the temperature 298.15 K 81

Bessergenev, V. G.; Kovalevskaya, Yu. A.; Paukov, I. E.; Starikov, M. A.; Oppermann, H.; Reichelt, W. Thermodynamic properties of $MnMnO_4$ and $Mn_2Mo_3O_8$ . . 85

Smith, J. R.; Zanonato, P. L.; Choppin, G. R. An elevated-temperature titration calorimeter 99

NOTE

Ribeiro da Silva, M. A. V.; Matos, M. A. R.; Yang Meng-Yan; Pilcher, G. Enthalpy of formation of 4-hydroxypyridine 107

Advice to authors 109

Number 2, *February* 1992

Pardo, J.; Rodríguez, V.; López, M. C.; Royo, F. M.; Urieta, J. S. Excess molar volumes V_m^E of (an alkanol + another alkanol) or (hexane + an alkanol) or (an alkanediol + an alkanol or another alkanediol) at the temperature 303.15 K 113

Souza, M. J.; Jiménez, E.; Legido, J. L.; Fernández, J.; Pérez Martell, E.; Paz Andrade, M. I. Excess molar volumes at the temperature 298.15 K of $\{x_1C_2H_5CO_2(CH_2)_2CH_3 + x_2Cl(CH_2)_5CH_3 + (1-x_1-x_2)c\text{-}C_6H_{12}\}$, $\{x_1C_2H_5CO_2(CH_2)_2CH_3 + x_2Cl(CH_2)_5CH_3 + (1-x_1-x_2)CH_3(CH_2)_4CH_3\}$, and $\{x_1C_2H_5CO_2(CH_2)_2CH_3 + x_2c\text{-}C_6H_{12} + (1-x_1-x_2)CH_3(CH_2)_4CH_3\}$ . . 119

Wieczorek, S. A. Thermodynamics of aqueous mixtures of polyoxyethene glycol monoethers. II. Excess molar volumes of (*n*-hexyl trioxyethene glycol monoether + water) 129

Huntelaar, M. E.; Cordfunke, E. H. P.; Ouweltjes, W. The standard molar enthalpies of formation of $SrSiO_3$(s) and Sr_2SiO_4(s). 139

Arda, N.; Sayar, A. A. (Liquid + liquid) equilibria of (water + tetrahydrofuran + 1-methylcyclohexanol) and of (water + tetrahydrofuran + 2-methylbutyl ethanoate) at the temperature (293.16 ± 0.30) K and pressure (101.325 ± 0.070) kPa 145

Kar, M.; Lenz, T. G.; Vaughan, J. D. Thermodynamic measurements for the Diels-Alder adduct of anthracene and maleic anhydride 151

Siu, W. W. Y.; Wong, T. Y. H.; Lai, J. T. W.; Chong, A.; Koga, Y. Isotope effects on the excess partial molar enthalpies of $\{xCH_3(CH_2)_3O(CH_2)_2O$(H or D) + $(1-x)(H_2O$ or $D_2O)\}$. 159

Sorai, M.; Kaji, K.; Kaneko, Y. An automated adiabatic calorimeter for the temperature range 13 K to 530 K. The heat capacities of benzoic acid from 15 K to 305 K and of synthetic sapphire from 60 K to 505 K 167

Saluja, P. P. S.; Lemire, R. J.; LeBlanc, J. C. High-temperature thermodynamics of aqueous alkali-metal salts 181

López, M.; Legido, J. L.; Romaní, L.; Carballo, E.; Pérez Martell, E.; Jiménez, E.; Paz Andrade, M. I. Excess molar enthalpies for (propan-1-ol + pentan-3-one + hexane) at the temperature 298.15 K 205

Acree, W. E., Jr.; Tucker, S. A.; Pilcher, G. Enthalpies of combustion of 1,4-dicyanobenzene di-*N*-oxide and 1,4-dicyanobenzene: at the mean dissociation enthalpy of the (N–O) bonds 213

Abboud, J.-L. M.; Jiménez, P.; Roux, M. V.; Turrión, C.; López-Mardomingo, C.; Sanz, G. Structural effects on the thermochemical properties of carbonyl compounds. II. Enthalpies of combustion, vapour pressures and enthalpies of sublimation, and standard enthalpies of formation in the gaseous phase, of 1-adamantyl methyl ketone and of 1,1′-diadamantyl ketone. 217

Number 3, *March* 1992

Konkova, T. S.; Matyushin, Ju. N.; Greenberg, J. H.; Lazarev, V. B.; Shapligin, I. S. Thermochemistry of double oxides in $Y_aBa_bCu_cO_d$ 225

Callanan, J. E.; McDermott, K. M.; Weir, R. D.; Westrum, E. F., Jr. Comparison of heat capacities measured by adiabatic calorimetry and by scanning calorimetry: thermodynamic properties of 9-methylcarbazole at temperatures between 4 K and 345 K 233

Steele, W. V.; Chirico, R. D.; Knipmeyer, S. E.; Nguyen, A. The thermodynamic properties of 9-methylcarbazole and of 1,2,3,4-tetrahydro-9-methylcarbazole . 245

Asou, M.; Terai, T.; Takahashi, Y. High-temperature enthalpy-increment measurements and derived heat capacity of lithium orthosilicate (Li_4SiO_4) at temperatures from 300 K to 1000 K 273

Asou, M.; Terai, T.; Takahashi, Y. An improved method of heat-capacity derivation from experimental enthalpies 281

Miyanaga, S.; Tamura, K.; Murakami, S. Excess enthalpies and excess isobaric heat capacities of (2,2,2-trifluoroethan-1-ol + benzene), (benzene + dimethyl-sulfoxide), and (2,2,2-trifluoroethan-1-ol + dimethylsulfoxide), at the temperature 298.15 K . 291

Barczyńska, J.; Bartel, L.; Mokrzan, J.; Taniewska-Osińska, S. Standard enthalpies of solution of zinc dichloride and of cadmium dichloride in (methanol + water) at the temperature 298.15 K 297

Ewing, M. B.; Goodwin, A. R. H. Thermophysical properties of alkanes from speeds of sound determined using a spherical resonator. 5. 2-Methylbutane at temperatures in the range 260 K to 320 K and pressures in the range 2.8 kPa to 80.9 kPa . 301

Holmes, H. F.; Mesmer, R. E. Isopiestic studies of H_2SO_4(aq) at elevated temperatures. Thermodynamic properties. 317

Cordfunke, E. H. P.; Konings, R. J. M. The vapour pressure of WCl_6. 329

Svoboda, V.; Kubeš, V.; Basařová, P. Enthalpies of vaporization and cohesive energies of hexan-2-one, 2-methylpentan-4-one, 2,2-dimethylbutan-3-one, 2,6-dimethylheptan-4-one, and cyclohexanone 333

Number 4, *April* 1992

Fuger, J. Transuranium-element thermochemistry: a look into the past – a glimpse into the future. (The 39th Huffman Memorial Award Lecture of the Calorimetry Conference delivered on 29 July 1991 at its 46th annual Conference held at Northern Illinois University, DeKalb, IL, U.S.A.) 337

Nguyen, S. N.; Silva, R. J.; Weed, H. C.; Andrews, J. E., Jr. Standard Gibbs free energies of formation at the temperature 303.15 K of four uranyl silicates: soddyite, uranophane, sodium boltwoodite, and sodium weeksite 359

Fernández, D. P.; Fernández-Prini, R. Solubility of solid iodine in near-critical fluid xenon 377

Walther, D.; Platzer, B.; Maurer, G. High-pressure (vapour + liquid) equilibria of (carbon dioxide + methylbenzene or 1,2-dimethylbenzene or 1,3-dimethylbenzene or 1,4-dimethylbenzene) at temperatures between 313 K and 393 K and pressures up to 17.3 MPa 387

Ribeiro da Silva, M. A. V.; Reis, A. M. M. V. Enthalpies of formation of crystalline dialkyldithiocarbamate complexes with iron(III), cobalt(III), chromium(III), and manganese(III): the mean M–S bond-dissociation enthalpies 401

Tillner-Roth, R.; Baehr, H. D. Burnett measurements and correlation of gas-phase (p, ρ, T) of 1,1,1,2-tetrafluoroethane (R 134a) and of 1,1-difluoroethane (R 152a) . 413

Roy, R. N.; Roy, L. N.; Ingle, A. L.; Davis, W. B.; Larkin, D. W.; Johnson, D. A.; Hnath, E. J.; Millero, F. J. The study of activity coefficients of (hydrogen bromide + strontium bromide)(aq) using Pitzer's formalism 425

Dias, A. R.; Minas da Piedade, M. E.; Martinho Simões, J. A.; Simoni, J. A.; Teixeira, C.; Diogo, H. P.; Yang, M.-Y.; Pilcher, G. Enthalpies of formation of *cis*-azobenzene and *trans*-azobenzene 439

Number 5, *May* 1992

Alonso, R.; Corrales, J. A.; Dominguez, A.; Tojo, J. Excess molar enthalpies of (methylcyclohexane + an alkanol) at the temperature 313.15 K. IV. General discussion of results 449

Zielkiewicz, J. (Vapour + liquid) equilibria in (propan-1-ol + heptane + octane) at the temperature 313.15 K 455

Gracia, M.; Sánchez, F.; Pérez, P.; Valero, J.; Gutiérrez Losa, C. Vapour pressures of (butan-1-ol + hexane) at temperatures between 283.10 K and 323.12 K . . 463

Quadri, S. K.; Kudchadker, A. P. The critical temperatures and critical pressures of some examples of (a polar substance + a non-polar substance) and of (a polar substance + another polar substance) 473

Wormald, C. J.; Smith, G. R. Excess enthalpies of (carbon dioxide + pentane or hexane or heptane or cyclohexane or benzene)(g) at temperatures from 432.2 K to 573.2 K and pressures up to 2.71 MPa 481

Wormald, C. J.; Yerlett, T. K. Molar enthalpy increments for $\{0.5(CH_3)_2CO + 0.5C_6H_6\}$ at temperatures up to 573.2 K and pressures up to 9.75 MPa 493

Steele, W. V.; Chirico, R. D.; Knipmeyer, S. E.; Nguyen, A. The thermodynamic properties of benzothiazole and benzoxazole 499

Ewing, M. B.; Goodwin, A. R. H. An apparatus based on a spherical resonator for measuring the speed of sound in gases at high pressures. Results for argon at temperatures between 255 K and 300 K and at pressures up to 7 MPa . . 531

Uosaki, Y.; Hamaguchi, T.; Moriyoshi, T. Excess molar volumes of (1,1,3,3-tetramethylurea + an alkan-1-ol) at the temperature 298.15 K 549

Svoboda, V.; Kubeš, V.; Basařová, P. Enthalpies of vaporization and cohesive energies of 1,1,2,2-tetrachloro-1,2-difluoroethane, 1,2-dibromoethane, 1-bromo-2-chloroethane, 1,3-dibromopropane, and 1,4-dibromo-2,3-dichloro-1,1,2,3,4,4-hexafluorobutane 555

NOTE

Uosaki, Y.; Kitaura, S.; Moriyoshi, T. Compressions of 4-methyl-1,3-dioxolan-2-one and of some alkanols at pressures up to 200 MPa and at the temperature 298.15 K . 559

Number 6, *June* 1992

Rogers, D. W.; Dejroongruang, K.; Samuel, S. D.; Fang, Wei; Zhao, Yeping Enthalpies of hydrogenation of the octenes and the methylheptenes 561

Callanan, J. E.; Weir, R. D.; Westrum, E. F., Jr. Thermodynamics of ammonium halogenated tellurides. I. Heat capacity and thermodynamic properties of $(NH_4)_2TeCl_6$ at temperatures from 5 K to 350 K 567

Vidavskii, L. M.; Korotkevich, I. I. The standard molar enthalpy of formation of heazlewoodite Ni_3S_2 determined by its synthesis in a calorimeter . . . 577

Ribeiro da Silva, M. A. V.; Ferrão, M. L. C. C. H.; Monte, M. J. S.; Esteves da Silva, R. M. G.; Ribeiro, J. C. Standard enthalpies of formation of 3-methylpentane-2,4-dione and of bis(3-methylpentane-2,4-dionato)copper(II): the mean (Cu O) bond-dissociation enthalpy 585

Ribeiro da Silva, M. A. V.; Ferrão, M. L. C. C. H.; Marques, R. M. C.; Lima, J. M. T. Standard enthalpies of formation of tetrakis(pentane-2,4-dionato)zirconium(IV) and tetrakis(1,1,1-trifluoropentane-2,4-dionato)zirconium(IV): the mean (Zr–O) bond-dissociation enthalpies 595

Jackowski, A. W. The effect of polymorphic forms of 4-(*trans*-4′-*n*-butylcyclohexyl) benzene isothiocyanate on the phase equilibria in *n*-pentane solutions . . . 601

Nagata, I.; Ozaki, S.; Myohen, K. Excess molar enthalpies for (acetonitrile + butan-2-one), (acetonitrile + trichloromethane + butan-2-one), and (trichloromethane + propan-2-one + benzene) 607

Nagata, I.; Tamura, K. Excess enthalpies of (aniline + acetonitrile or benzene) and of (aniline + acetonitrile + benzene) at the temperature 298.15 K 613

Apelblat, A. The vapour pressures of water over saturated aqueous solutions of barium chloride, magnesium nitrate, calcium nitrate, potassium carbonate, and zinc sulfate, at temperatures from 283 K to 313 K 619

Callanan, J. E.; Hope, G. A.; Weir, R. D.; Westrum, E. F., Jr. Thermodynamic properties of tungsten ditelluride (WTe_2). 1. The preparation and low-temperature heat capacity at temperatures from 6 K to 326 K 627

O'Hare, P. A. G.; Hope, G. A. Thermodynamic properties of tungsten ditelluride (WTe_2). II. Standard molar enthalpy of formation at the temperature 298.15 K . 639

Hongo, M.; Kusunoki, M.; Kumagae, Y.; Arai, Y. (Vapour + liquid) equilibria of (propan-2-ol + propan-1-ol) and of (propan-2-ol + propan-1-ol + calcium chloride) at the temperature 298.15 K 649

Ribeiro da Silva, M. A. V.; Reis, A. M. M. V.; Monte, M. J. S.; Bártolo, M. M. S. S. F.; Rodrigues, J. A. R. G. O. Enthalpy of combustion, vapour pressures, and enthalpy of sublimation of 3-nitrophenol 653

Callanan, J. E. Weir, R. D.; Westrum, E. F., Jr. Thermodynamics of ammonium halogenated tellurides. II. Heat capacity, phase transitions, and thermodynamic properties of $(ND_4)_2TeCl_6$ at temperatures from 5 K to 325 K 661

Number 7, *July* 1992

Lebedev, B. V.; Kulagina, T. G.; Sheiman, M. S.; Nistratov, V. P. Thermodynamics of the complexes $(CH_3)_2Cd \cdot Se(CH_3)_2$ and $(CH_3)_2Cd \cdot Te(CH_3)_2$ in the temperature range from 0 to 310 K. 673

Händel, G.; Kleinrahm, R.; Wagner, W. Measurements of the (pressure, density, temperature) relation of methane in the homogeneous gas and liquid regions in the temperature range from 100 K to 260 K and at pressures up to 8 MPa . . 685

Händel, G.; Kleinrahm, R.; Wagner, W. Measurements of the (pressure, density, temperature) relations of dichlorodifluoromethane (R12) and of chlorodifluoromethane (R22) in parts of the homogeneous gas and liquid regions and on the coexistence curve 697

Ribeiro da Silva, M. A. V.; Monte, M. J. S. Vapour pressures and enthalpies of sublimation of six halogen-substituted 8-hydroxyquinolines. 715

Cordfunke, E. H. P.; Konings, R. J. M.; Meyssen, S. R. M. Vapour pressures of some caesium compounds. II. Cs_2MoO_4 and Cs_2RuO_4 725

Banipal, T. S.; Garg, S. K.; Ahluwalia, J. C. Densities of some higher alkan-1-oic acids at temperatures from 343.15 K to 373.15 K and at pressures up to 9 MPa . . 729

Zaitsev, A. I.; Litvina, A. D.; Mogutnov, B. M. Thermodynamic properties of the intermediate phases of $\{xCaF_2 + ySiO_2 + (1-x-y)CaO\}$ 737

Lebedev, B. V.; Kulagina, T. G. Thermodynamic properties of 1,2-difluoro-1,2,2-trichloroethane and of 1,2-difluoro-2,2-dichloroethane at temperatures from 0 to 330 K 751

Wirbser, H.; Bräuning, G.; Gürtner, J.; Ernst, G. Flow-calorimetric specific heat capacities and Joule-Thomson coefficients of CF_3CHFCF_3 (R227)—a new working fluid in energy technics—at pressures up to 15 MPa and temperatures between 253 K and 423 K 761

Brumleve, T. R.; Walston, S. R.; Mucklejohn, S. A.; O'Brien, N. W. The preparation, vapour pressure, and thermodynamic properties of tin(II) bromide 773

NOTE

Wirbser, H.; Bräuning, G.; Ernst, G. Flow-calorimetric specific heat capacities of the refrigerants $CFCl_3$ (R11) and $CF_2ClCFCl_2$ (R113) at pressures between 0.6 MPa and 30 MPa and temperatures between 288.15 K and 503.15 K 783

Number 8, *August* 1992

Kabata, Y.; Yamaguchi, S.; Takada, M.; Uematsu, M. Densities of 2,2,2-trifluoroethanol in the temperature range from 310 K to 420 K. II. Compressed-liquid densities at pressures up to 200 MPa 785

Uosaki, Y.; Iwama, F.; Moriyoshi, T. Compressions of (water + formamide or *N*,*N*-dimethylformamide) at pressures up to 150 MPa and at the temperature 298.15 K 797

López, M.; Fernández, J.; Sarmiento, F.; Legido, J. L.; Romaní, L.; Jiménez, E.; Paz Andrade, M. I. Excess molar enthalpies at the temperature 298.15 K of (an *n*-alkyl formate + an *n*-alkanol). IV. $\{xHCO_2(CH_2)_iCH_3 + (1-x)C_jH_{2j+1}OH\}$, ($i = 0$ to 3 and $j = 1$ and 2) 809

Cordfunke, E. H. P.; Laan, R. R. van der; Westrum, E. F., Jr. The thermochemical and thermophysical properties of Cs_2RuO_4 and Cs_2MnO_4 at temperatures from 5 K to 1000 K 815

Chao, J.-P.; Zhang, F.-Q.; Dai, M. Studies of thermodynamic properties of binary mixtures containing an alcohol. XVIII. Excess molar enthalpies of each of (one of the four butanols + trichloromethane or 1,2-dichloroethane) 823

Holmes, H. F.; Mesmer, R. E. Isopiestic studies of aqueous solutions at elevated temperatures. XI. $\{(1-y)CsCl + yBaCl_2\}$(aq) 829

Gracia, M.; Sánchez, F.; Pérez, P.; Valero, J.; Gutiérrez Losa, C. Vapour pressures of $\{xC_4H_9OH + (1-x)C_8H_{18}\}$(l) at temperatures between 283.16 K and 323.18 K . 843

Shen, S.; Wang, Y.; Shi, J.; Benson, G. C.; Lu, B. C.-Y. Excess enthalpies of (1,2-dichloroethane + an alkan-1-ol) and of (3-oxa-pentane-1,5-diol + an alkan-1-ol) 851

Kemp, W. J. M. van der; Oonk, H. A. J.; Bergman, C.; Chastel, R. Determination of the excess molar Gibbs energy of $\{(1-x)BaO + xSrO\}(s)$ by means of Knudsen-effusion mass spectrometry 857

Fujiwara, T.; Inaba, A.; Atake, T.; Chihara, H. Thermodynamic properties of deuterated hexamethylbenzene and of its solid solutions with the hydrogenated analog. A large isotope effect on the phase transition at the temperature 117 K 863

Kozyro, A. A.; Kabo, G. J.; Krouk, V. S.; Sheiman, M. S.; Yursha, I. A.; Simirsky, V. V.; Krasulin, A. P.; Sevruk, V. M.; Gogolinsky, V. I. Thermodynamic properties of cyclohexanone oxime 883

Number 9, *September* 1992

Shukla, N. K.; Prasad, R.; Roy, K. N.; Sood, D. D. Thermochemistry of lithium chromate Li_2CrO_4(cr) and lithium molybdate Li_2MoO_4(cr). 897

Prieto, G.; Nogueira, P.; Bravo, R.; Sarmiento, F.; Paz Andrade, M. I. Excess enthalpies and excess volumes of (pentan-2-one + an *n*-alkan-1-ol) at the temperature 298.15 K 905

Grønvold, F.; Stølen, S. Thermodynamics of iron sulfides. II. Heat capacity and thermodynamic properties of FeS and of $Fe_{0.875}S$ at temperatures from 298.15 K to 1000 K, of $Fe_{0.98}S$ from 298.15 K to 800 K, and of $Fe_{0.89}S$ from 298.15 K to about 650 K. Thermodynamics of formation 913

Todorović, M.; Ninković, R. Osmotic and activity coefficients of $\{xKNO_3 + (1-x)Mg(NO_3)_2\}$(aq) at the temperature 298.15 K 937

Wormald, C. J.; Hodgetts, R. W.; Smith, G. R. Excess enthalpies of (carbon dioxide + cyclohexane)(g) and of (carbon dioxide + benzene)(g) at the pressure 101.3 kPa over the temperature range 354 K to 399 K 943

Gilgen, R.; Kleinrahm, R.; Wagner, W. Measurements of the (pressure, density, temperature) relation of sulphur hexafluoride (SF_6) in the homogeneous region at temperatures from 321.15 K to 333.15 K and at pressures up to 8 MPa and on the coexistence curve at temperatures from 288.15 K to 315.15 K 953

Woolley, A. T.; Sipowska, J. T.; Ott, J. B.; Izatt, R. M. Excess enthalpies for (butane + acetonitrile) at the temperatures (298.15, 323.15, and 348.15) K and at the pressures (5, 10, and 15) MPa 965

Trusler, J. P. M.; Zarari, M. The speed of sound and derived thermodynamic properties of methane at temperatures between 275 K and 375 K and pressures up to 10 MPa. 973

Calado, J. C. G.; Lopes, J. N. C.; Rebelo, L. P. N. Vapour pressure of trideuterioammonia 993

Callanan, J. E.; Weir, R. D.; Westrum, E. F., Jr. Thermodynamics of ammonium halopallidates. I. Heat capacity and thermodynamic properties of $(NH_4)_2PdCl_6$ at temperatures from 6 K to 350 K. 1001

Number 10, *October* 1992

O'Hare, P. A. G.; Susman, S.; Volin, K. J.; Rowland, S. C. Combustion calorimetry of rhombohedral sulfur in fluorine: a question of impurities. The standard molar enthalpies of formation of SF_6(g) and SO_4^{2-}(aq) at the temperature 298.15 K 1009

Kabata, Y.; Yamaguchi, S.; Takada, M.; Uematsu, M. Densities of 2,2,2-trifluoroethanol in the temperature range from 310 K to 420 K. I. Saturated-liquid densities. . 1019

Jain, D. V. S.; Chadha, R.; Dhar, N. S. Excess molar enthalpies of (benzene or methylbenzene or ethylbenzene + 2-methylethylbenzene) at the temperatures 298.15 K, 308.15 K, and 318.15 K 1027

Kato, Y.; Asano, M.; Harada, T.; Mizutani, Y. Mass-spectrometric study of the thermodynamic properties of $\{xRbBO_2 + (1-x)CsBO_2\}$ 1033

Zaitsev, A. I.; Litvina, A. D.; Mogutnov, B. M. Thermodynamic properties of $\{xCaF_2 + ySiO_2 + (1-x-y)CaO\}$(l). 1. Experimental 1039

Gruszkiewicz, M. S.; Sipowska, J. T.; Ott, J. B.; Graham, R. C. Excess enthalpies for (butane + chloromethane) at the temperatures (298.15, 323.15, and 348.15) K and pressures (5 and 15) MPa and at the temperature 293.15 K and pressure 15 MPa 1057

Salvi-Narkhede, M.; Wang, B.-H.; Adcock, J. L.; Van Hook, W. A. Vapor pressures, liquid molar volumes, vapor non-ideality, and critical properties of some partially fluorinated ethers ($CF_3OCF_2CF_2H$, CF_3OCF_2H, and CF_3OCH_3), some perfluoroethers ($CF_3OCF_2OCF_3$, *c*-$CF_2OCF_2OCF_2$, and *c*-$CF_2CF_2CF_2O$), and of CHF_2Br and CF_3CFHCF_3 1065

Miyanaga, S.; Tamura, K.; Murakami, S. Excess molar volumes, isentropic and isothermal compressibilities, and isochoric heat capacities of (acetonitrile + benzene), (benzene + dimethylformamide), and (acetonitrile + dimethylformamide) at the temperature 298.15 K 1077

Sipowska, J. T.; Ott, J. B.; Woolley, A. T.; Izatt, R. M. Excess enthalpies for (butane + methanol) at the temperatures (298.15, 323.15, and 348.15) K and the pressures (5 and 15) MPa 1087

Naumowicz-Węglińska, E.; Wóycicki, W. Excess enthalpies of gaseous mixtures containing ammonia. 3. (Ammonia + ethene) 1095

Huntelaar, M. E.; Cordfunke, E. H. P.; Ouweltjes, W. The standard molar enthalpies of formation of $BaSiO_3$(s) and Ba_2SiO_4(s) 1099

Vis, M. G. M. van der; Cordfunke, E. H. P.; Konings, R. J. M.; Berg, G. J. K. van den; Miltenburg, J. C. van Tetraethoxysilane, $Si(OC_2H_5)_4$: heat capacity and thermodynamic properties at temperatures from 0 to 440 K 1103

Liu, Y.-Q.; Maurer, G. Flow calorimeter for measuring enthalpy changes during the partial vaporization of mixtures 1109

NOTE

Powell, B. M.; Weir, R. D. Thermodynamic properties of ammonium halogen stannates. II. Neutron diffraction and heat capacity of $(ND_4)_2SnCl_6$ 1119

Number 11, *November* 1992

Ortega, J. Excess molar enthalpies at the temperature 298.15 K of (a methyl *n*-alkanoate + pentane or heptane) 1121

Zaitsev, A. I.; Zemchenko, M. A.; Litvina, A. D.; Mogutnov, B. M. Thermodynamic properties of the alloys $\{xBa + (1-x)P\}$ 1127

Wang, L.; Benson, G. C.; Lu, B. C.-Y. Excess enthalpies of (ethanol + hexane + decane or dodecane) at the temperature 298.15 K 1135

Jiménez, P.; Roux, M. V.; Turrión, C. Thermochemical properties of *N*-heterocyclic compounds. IV. Enthalpies of combustion, vapour pressures and enthalpies of sublimation, and enthalpies of formation of 2-methylimidazole and 2-ethylimidazole 1145

Boyes, S. J.; Ewing, M. B.; Goodwin, A. R. H. Heat capacities and second virial coefficients for gaseous methanol determined from speed-of-sound measurements at temperatures between 280 K and 360 K and pressures from 1.03 kPa to 80.5 kPa 1151

Tomas, G.; Artal, M.; Otín, S. Excess enthalpies of (*N*,*N*-dimethylformamide or *N*,*N*-dimethylacetamide + hexane or benzene or toluene or *p*-xylene or mesitylene) 1167

Sharma, S. C.; Kaur, R.; Singh, J. Excess molar volumes of (methyl 1,1-dimethylethyl ether + cyclohexane or benzene or toluene or ethyl benzene or propyl benzene or *o*-xylene or *m*-xylene or *p*-xylene) at the temperature 303.15 K 1171

Nagata, I.; Miyazaki, K. Excess enthalpies of (aniline + propan-2-ol) and of (aniline + propan-2-ol + benzene) at the temperature 298.15 K 1175

Konings, R. J. M.; Booij, A. S. High-temperature i.r. spectroscopy of the vapour over $AlCl_3$. The thermodynamic properties of $AlCl_3(g)$ and $Al_2Cl_6(g)$ 1181

Ohta, Y.; Matsuo, T.; Suga, H. Heat capacity and phase transitions of scandium ethanoate 1189

Keller, M.; Heintz, A.; Lichtenthaler, R. N. Excess molar enthalpies of (tetrahydrofuran + methanol + water) at the temperature 298.15 K and pressure 0.1 MPa 1197

Malhotra, R.; Woolf, L. A. Thermodynamic properties of butan-2-one at temperatures from 278 K to 338 K and pressures from 0.1 MPa to 280 MPa; predictions for higher ketones 1207

Ribeiro da Silva, M. A. V.; Monte, M. J. S. Vapour pressures and standard molar enthalpies of sublimation of four crystalline β-diketones 1219

Letcher, T. M.; Mercer-Chalmers, J.; Bean, B. A. J.; Battino, R. Excess molar volumes and excess molar enthalpies of $[x\{0.5CH_3(CH_2)_{k-2}CH_3 + 0.5CH_3(CH_2)_{l-2}CH_3\} + (1-x)CH_3(CH_2)_{m-2}CH_3]$ at the temperature 298.15 K . 1229

Number 12, *December* 1992

Avedis, G.; Roux, A. H.; Grolier, J.-P. E. Excess molar enthalpies of (tetrachloromethane + an *n*-alkyl ester) at the temperature 298.15 K . . . 1233

Gracia, M.; Baños, I.; Pérez, P.; Valero, J.; Gutiérrez Losa, C. Thermodynamics of the partially miscible $\{x(CH_3)_2C(OH)CH_2COCH_3 + (1-x)CH_3(CH_2)_4CH_3\}$ at temperatures between 288.15 K and 318.15 K 1237

Gilgen, R.; Kleinrahm, R.; Wagner, W. Supplementary measurements of the (pressure, density, temperature) relation of carbon dioxide in the homogeneous region at temperatures from 220 K to 360 K and pressures up to 13 MPa 1243

Cordfunke, E. H. P.; Laan, R. R. van der; Wyers, G. P.; Miltenburg, J. C. van The heat capacities and derived thermophysical properties of Li_2ZrO_3 and Li_8ZrO_6 at temperatures from 0 to 1000 K 1251

Ewing, M. B.; Goodwin, A. R. H. Speeds of sound, perfect-gas heat capacities, and acoustic virial coefficients for methane determined using a spherical resonator at temperatures between 255 K and 300 K and pressures in the range 171 kPa to 7.1 MPa 1257

Korobov, M. V.; McCreary, J. C.; Kematick, R. J.; Myers, C. E. The ion-current-ratio method in Knudsen-cell mass spectrometry: binary mixtures with stoichiometric intermediate phases 1275

Aguiar Ricardo, A.; Barreiros, S. F.; Nunes da Ponte, M.; Albuquerque, G. M. N.; Calado, J. C. G. (p, V_m, T) of (0.476Ar + 0.524N_2)(l) and the calculation of thermodynamic properties of liquid air 1281

Ribeiro da Silva, M. A. V.; Ferrão, M. L. C. C. H.; Esteves da Silva, R. M. G. Standard molar enthalpy of formation of tris(3-methylpentane-2,4-dionato)iron(III): the mean (Fe–O) bond-dissociation enthalpy 1293

Abboud, J.-L. M.; Jiménez, P.; Roux, M. V.; Turrión, C.; López-Mardomingo, C. Structural effects on the thermochemical properties of organic compounds. III. Enthalpies of combustion, vapour pressures and enthalpies of sublimation, and standard enthalpies of formation in the gaseous phase, of adamantane-1-carboxylic acid methyl ester and of adamantane-1-carbonitrile 1299

Wang, L.; Benson, G. C.; Lu, B. C.-Y. Excess enthalpies of {(ethanol or propan-1-ol or methyl 1,1-dimethylethyl ether) + (2,3-dimethylbutane or 2,2,4-trimethylpentane)} . 1305

Diaz, E. L.; Domalski, E. S.; Colbert, J. C. Enthalpies of combustion of glycylglycine and DL-alanyl-DL-alanine 1311

Zielkiewicz, J. (Vapour + liquid) equilibria in (2-methylpyridine + benzene or ethyl ethanoate or chlorobenzene) at the temperature 303.15 K 1319

O'Hare, P. A. G. Thermodynamic properties of silicides I. Calorimetric measurements of the specific energy of combustion in fluorine of hyperstoichiometric tungsten disilicide, and the derived standard molar enthalpy of formation $\Delta_f H^\circ_m(WSi_{2.060})$ at the temperature 298.15 K 1323

INDEX TO VOLUME 24 1333